



Braga Goncalves, I., Cornetti, L., Couperus, A., van Damme, C., & Mobley, K. (2017). Phylogeography of the snake pipefish, *Entelurus aequoreus* (Family: Syngnathidae) in the northeastern Atlantic Ocean. *Biological Journal of the Linnean Society*, [blx112].
<https://doi.org/10.1093/biolinnean/blx112>

Peer reviewed version

License (if available):
Unspecified

Link to published version (if available):
[10.1093/biolinnean/blx112](https://doi.org/10.1093/biolinnean/blx112)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://academic.oup.com/biolinnean/article/doi/10.1093/biolinnean/blx112/4555402/Phylogeography-of-the-snake-pipefish-Entelurus>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Original Article

Phylogeography of the snake pipefish, *Entelurus aequoreus* (Family: Syngnathidae) in the northeastern Atlantic Ocean

Ines Braga Goncalves¹, Luca Cornetti², Abraham S. Couperus³, Cindy J.G. van Damme³,
Kenyon B. Mobley⁴

¹*School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, Bristol, BS8 1TQ, UK*

²*Zoological Institute, University of Basel, Vesalgasse 1, CH-4051 Basel, Switzerland*

³*Wageningen Marine Research (IMARES), Postbus 68, 1970 AB IJmuiden, The Netherlands*

⁴*Department of Evolutionary Ecology, Max Planck Institute for Evolutionary Biology, August-Thienemann Str. 2, 24306, Plön, Germany*

Corresponding author: Kenyon Mobley

Email: mobley@evolbio.mpg.de

Phone: +49 4522.763.347

Running title: Phylogeography of the snake pipefish

ABSTRACT

The snake pipefish, *Entelurus aequoreus*, is a widespread marine species occurring in pelagic and coastal environments in the northeastern Atlantic Ocean. Recently, the snake pipefish underwent a short-lived, yet substantial, increase in abundance and range expansion into arctic waters. However, little is known about the species' population structure or if different ecotypes contributed to this outbreak. Specimens (n=178) were sampled from 25 locations from six regions spanning 1.9 million km². A fragment of the mitochondrial cytochrome *b* gene and control region was used to assess population structure and genetic diversity. Both loci showed high haplotype diversity (H_d) and low nucleotide diversity (π) over all sampled locations. A genetic signature of population expansion was evident through mismatch distributions and tests for recent population expansion (Fu's F_s , Tajima's D , and R_2). Effective population size analyses (Bayesian Skyline Plot) suggest an ancient expansion (50-100 thousand years ago). However, we found neither significant population differentiation (AMOVA) among regions, nor evidence of genetically distinct ecotypes. This lack of structure is likely due to a pelagic life style, fast development and long distance dispersal aided by ocean currents. Our work highlights the need for further research to better understand the recent outbreak and how this species may respond to future environmental challenges.

Keywords: Bayesian Skyline Plot, control region, cytochrome *b*, fish, life history, mitochondrial DNA, pelagic, population increase, population structure, range expansion

Introduction

The introduction of molecular techniques to the study of species distributions has greatly improved our knowledge regarding species' evolutionary history. Current genetic tools allow scientists to assess historical species distribution patterns, gene flow between populations, the identification of source populations, routes and patterns of invasion, and timing of range shifts, expansions as well as contractions (Hewitt, 1999; Schmitt, 2007; Moran & Alexander, 2014). For example, the assessment of levels of gene flow and genetic diversity between geographically separate populations is of particular interest for conservation practises because it can facilitate the identification of genetically independent populations or units, and guide conservation efforts (e.g., Bernard, Feldheim, Heithaus *et al.*, 2016). These studies highlight the importance of linking genetic data with species historical and geographical information to make biologically relevant interpretations. The discipline of phylogeography does exactly that by looking at geographical patterns of genetic diversity across populations or species over time (Avice, 2000).

In this study, we investigate the phylogeography of the snake pipefish, *Entelurus aequoreus* L. 1758. This species is a member of the family Syngnathidae, the pipefishes, seahorses and seadragons, a group characterized by its unique form of reproduction, male pregnancy (Dawson, 1986). Life history traits appear to shape the past and present geographical distributions of syngnathids (Mobley, Small & Jones, 2011). Syngnathids tend to be poor swimmers with small fins and armoured bodies, and most are strict habitat specialists, relying heavily on crypsis for survival (Vincent, Berglund & Ahnesjö, 1995). All species in this family produce free-living juveniles (Hasse, 1974; Mi, Kornienko & Drozdov, 1998; Monteiro, Almada & Vieira, 2003; Wetzel & Wourms, 2004) with short (Wilson & Vincent, 1998; Planas, Blanco, Chamorro *et al.*, 2012) or entirely absent pelagic dispersal phases (Silva, Monteiro, Almada *et al.*, 2006; Silva, Monteiro, Vieira *et al.*, 2006). Having a

short pelagic dispersal phase is known to significantly limit dispersal potential (Grantham, Eckert & Shanks, 2003). Among adults, limited seasonal vertical migrations can occur, with individuals of some species coming into warmer shallow waters for mating, and returning to deeper waters at the end of breeding season or during brooding (Vincent *et al.*, 1995; Monteiro, Berglund, Vieira *et al.*, 2006).

Previous studies on syngnathids have generally revealed relatively high genetic diversity and high population structuring, indicative of large effective population sizes and low dispersal ability (reviewed in Mobley *et al.*, 2011). Most northern-hemisphere temperate syngnathid species show evidence of population expansions towards northern regions after the end of the last glaciation period (circa 20 thousand years before present (ka); Woodall, Koldewey, Santos *et al.*, 2009; Mobley, Small, Jue *et al.*, 2010; Wilson & Eigenmann Veraguth, 2010; Woodall, Koldewey, Boehm *et al.*, 2015) following similar northerly expansion of suitable habitat, i.e., seagrass meadows (Olsen, Stam, Coyer *et al.*, 2004). Yet, their strong habitat dependency and limited dispersal capability is also reflected in their current geographical distributions that generally show high levels of population differentiation on relatively small geographical scales (e.g. Lourie, Green & Vincent, 2005; Wilson, Stiller & Rouse, 2016; Stiller, Wilson, Donnellan *et al.*, 2017) and strong indications of limited dispersal (e.g. Chenoweth, Hughes & Connolly, 2002; reviewed in Mobley *et al.*, 2011; Wilson & Orr, 2011). Exceptions to these patterns have been interpreted as a result of recent colonization events (Nickel & Cursons, 2012), assumed to occur via rafting, where individuals drift with floating marine vegetation on ocean currents (Teske, Hamilton, Palsbøll *et al.*, 2005; Fedrizzi, Stiassny, Boehm *et al.*, 2015).

From a phylogeographical perspective, the snake pipefish is an interesting species to study because of its wide geographic distribution, recent range expansion into polar waters, and dramatic fluctuations in abundance. Historically, this species inhabits a vast geographical

range in the temperate northeastern Atlantic, spanning from the European continental shelf to the west, southern Norway and Iceland in the north, to the Azores in the south and the Baltic Sea to the east (Dawson, 1986). Normally, the species is encountered throughout its range at low densities (Harris, Beare, Toresen *et al.*, 2007). However, between 2003 and 2007 the snake pipefish reappeared in large numbers in coastal areas where it had for decades been considered rare (e.g. northern Wadden Sea, Polte & Buschbaum, 2008), including brackish estuarine waters (e.g. the Severn Estuary, Henderson & Bird, 2010). During this time, the snake pipefish was caught in numbers several orders of magnitude higher than in catches prior to 2003 (Kloppmann & Ulloweit, 2007; van Damme & Couperus, 2008). The snake pipefish also underwent a geographical range expansion into the Barents and Greenland Seas by 2005 (Harris *et al.*, 2007; Rusyaev, Dolgov & Karamushko, 2007) and the first ever records of occurrence in Svalbard were reported by August 2006, representing a 15° latitudinal expansion northwards (approximately 1650km, Fleischer, Schaber & Piepenburg, 2007). After 2007, populations of the snake pipefish declined dramatically and returned to low levels of abundance throughout its geographic range (Heath, Neat, Pinnegar *et al.*, 2012). There is anecdotal evidence that this species has undergone mass mortality events off the European continental shelf in the Atlantic Ocean in 1885, 1887, and in the North Sea in 1911, although no satisfactory explanation for these events exists (Brongersma-Sanders, 1957). It is possible that these mass mortality events are an indication of previous population increases in snake pipefish abundance although data to corroborate this link are currently lacking.

The snake pipefish is also an interesting species to investigate phylogeographically because of its unique life history traits. Unlike most other syngnathids that are predominantly associated with benthic habitats, the snake pipefish is described primarily as an oceanic species displaying a pelagic lifestyle and is found in both coastal and oceanic waters to depths up to 100m (Dawson, 1986; Kloppmann & Ulloweit, 2007). Because this species does

not require benthic habitats to reproduce they may breed in open waters and offspring as well as adults may be transported and mixed by ocean currents.

Finally, the potential for cryptic species to contribute to the temporary population increase and range expansion of the snake pipefish has not yet been addressed. Previously, two species have been proposed for *E. aequoreus* over the years based on differences in body size, colour, position of the dorsal fin and in number of rays (Yarrel, 1839; Moreau, 1881; Fries, Ekström & Sundevall, 1895; Holt & Byrne, 1906; Dunker, 1915). However, these two species are currently not recognized (Dawson, 1986). Additional lines of investigation have suggested coastal and oceanic habitat-specific phenotypes, or ‘ecotypes’, based on morphology (van Damme & Couperus, 2008) or timing of breeding between coastal benthic populations (summer) and oceanic pelagic populations (spring) (Kloppmann & Ulloweit, 2007). Thus, the potential for ecotypes or cryptic species to exist within *E. aequoreus* needs to be resolved with molecular markers.

In this study, we investigate phylogeographic patterns in the snake pipefish throughout its contemporary distribution. Our specific goals are to assess population structure and historical patterns of population expansion over the geographical range of the snake pipefish. Further, we investigate whether molecular data support proposed coastal benthic and pelagic ecotypes or cryptic species. To achieve these goals, we used mitochondrial DNA (mtDNA, cytochrome *b* and control region) markers to investigate genetic differentiation and genetic variation in snake pipefish from 25 locations among six geographical regions spanning over 1.9 million km² of their range in the Northeastern Atlantic Ocean.

Materials and methods

Collections

A total of 237 snake pipefish were collected from 25 locations in the Northeastern Atlantic Ocean between 2003 and 2010 using a variety of capture methods (Table 1, Fig. 1, Supplementary file S1). We assigned samples to regions based on conventional naming schemes of local water bodies or coastal areas. These regions correspond to regions defined by the OSPAR Commission (Region I: Arctic Waters = Norwegian Sea; Region II: Greater North Sea = North Sea; Region III Celtic Seas = Continental Shelf; Region IV Bay of Biscay and Iberian Coast = Spanish Coast) with the exception that Skagerrak/Kattegat were analysed separately from the North Sea due to the potential influence of the Baltic Sea, and the French Coast was analysed separately as it lies on the boundary of North Sea and Celtic Seas regions.

DNA was extracted from a small piece of tail tissue (~1cm) using a Qiagen DNeasy kit from live, frozen or EtOH preserved fish. A portion of the mitochondrial cytochrome *b* gene and control region was amplified. We used primers L14725 (Pääbo, Thomas, Whitfield *et al.*, 1991) and H15926 (Wilson, Vincent, Ahnesjö *et al.*, 2001) to amplify the cytochrome *b* locus, and primers L15926 (Kocher, K., Meyer *et al.*, 1989) and H16498 (Meyer, Kocher, Basasibwaki *et al.*, 1990) to amplify the control region. Fragments were amplified via polymerase chain reaction (PCR) in 30µl volumes containing 3µl of 10X buffer, 1.8 µl of dNTPs (10µM of each dNTP), 3 µl of MgCl₂ (50mM), 1.5 µl of each primer (10µM), and 0.5 µl of Taq (5 units/µl; InviTaq, Stratech Biomedical, Birkenfeld, DE). The PCR thermal profile consisted of an initial denaturation at 95°C (2 min), followed by 35 cycles of 94°C (30 sec), reannealing temperature (30 sec), 72°C (90 sec), and a final extension at 72°C for 10 min. Reannealing temperature for cytochrome *b* was 48°C and control region was 56°C. PCR products were purified before sequencing with Illustra™ ExoStar (GE Healthcare, Buckinghamshire, UK) using 5µl of PCR product and 2µl of ExoStar. PCR products were

sequenced with the forward and reverse primers using an ABI 3100 DNA Analyzer (Applied Biosystems, Foster City, CA, USA).

Contigs were created using forward and reverse sequences using CODONCODE ALIGNER v. 5.1.5 (Codon Code Corporation, Centerville, MA, USA) for each individual and aligned for each locus using MUSCLE (Edgar, 2004). Sequences were verified by eye and trimmed. Unique haplotype sequences were identified using ‘DNA to haplotype collapse and converter’ in FaBox v. 1.41 (<http://birc.au.dk/fabox>) and deposited in GenBank (accession numbers #: cytochrome *b*, KY857646 - KY857823; control region, KY965149 - KY965308).

Genetic analyses

Relationships between mitochondrial haplotypes were analysed for each locus independently to assess whether population structuring exists among the six regions combined over all time periods. Standard haplotype (h) and nucleotide (π) diversity statistics (Nei, 1987) were calculated for each region using DNASP 5.10.01 (Librado & Rozas, 2009). Mismatch distributions were investigated in both loci independently, and evidence for recent population expansion was tested using Tajima’s D test (Tajima, 1989), Fu’s F_s test (Fu, 1997) and the Ramos-Onsins and Rozas’s R_2 statistic (Ramos-Onsins & Rozas, 2002), recommended for small sample sizes (Ramírez-Soriano, Ramos-Onsins, Rozas *et al.*, 2008). Tajima’s D , F_s and R_2 were calculated using the total number of mutations, excluding sites with alignment gaps or missing data, and significance was ascertained using coalescent simulations with 1000 replicates as implemented by DNASP. Population expansion under the constant size growth model was used to estimate R_2 .

To visualize the relationship between mitochondrial haplotypes from different regions, a haplotype network was constructed using HAPLOVIEWER (Salzburger, Ewing &

von Haeseler, 2011) based on maximum likelihood trees drawn in DNAML in PHYLIP v3.695 (Felsenstein, 1989) for both loci independently. To test for significant population subdivision among individual collections, we conducted pairwise ϕ_{ST} tests for cytochrome *b* and control region sequences using ARLEQUIN v3.5.2.1 (Excoffier, Smouse & Quattro, 1992; Excoffier, Laval & Schneider, 2005). Significance for pairwise differences was ascertained using an exact test with 100,000 permutations. An Analysis of Molecular Variance (AMOVA) was used to test the proportion of genetic differentiation within and between regions using ARLEQUIN for both loci independently pooled across years. AMOVA was also used to test whether there is support for genetic differentiation between pelagic and coastal benthic populations. Only locations with a minimum of five sequenced individuals were included in pairwise ϕ_{ST} and AMOVA analyses, except for the three Norwegian Sea individuals that were included in the cytochrome *b* dataset since they represented a unique location. Significance of AMOVAs was determined using 99,999 permutations as implemented in ARLEQUIN.

We analysed the potential for fluctuations in effective population size using Bayesian Skyline Plot (BSP, Drummond, Rambaut, Shapiro *et al.*, 2005), a coalescent-based method implemented in BEAST 1.8.4 (Drummond & Rambaut, 2007). We first estimated the best-fit model of nucleotide substitution for each gene using JModelTest 2.0 (Darriba, Taboada, Doallo *et al.*, 2012) on all samples. We selected the best-fit models according to the Bayesian Information Criterion, which were HKY+I+G and HKY+I for cytochrome *b* and the control region, respectively. We then ran the BSP analysis on the concatenation of the two mitochondrial loci using samples that successfully amplified both cytochrome *b* and the control region (n=140) using specific substitution models and mutation rates for each partition. We chose a mutation rate of 1×10^{-8} substitutions per nucleotide site per year for cytochrome *b* and 5×10^{-8} substitutions per nucleotide site per year for the control region

based on recommendations by Bowen, Muss, Rocha *et al.* (2006) and Mobley *et al.* (2010) assuming a constant mutation rate. We assumed a generation time of one generation per year based on unimodal distributions of body size in juveniles caught in plankton tows (Kirby, Johns & Lindley, 2006). Adults collected within a year show a bimodal distribution in body size but this may be accounted for by sexual dimorphism in body length (van Damme & Couperus, 2008) also suggesting one reproductive cycle per year.

The BSP analysis consisted of 1×10^8 generations; parameters were sampled every 10,000 generations of which 10% was discarded as burn-in. In order to check the analysis performance (i.e., the convergence of parameters by visually checking the effective sample size (ESS>200) values), we used TRACER 1.6 (Rambaut, Suchard, Xie *et al.*, 2014).

Ethical statement

The present study was conducted in accordance to local and European Union law and no permit was needed for collecting fish. The study did not involve any endangered or protected species.

Results

Molecular diversity

We resolved 900 bp of the cytochrome *b* gene from 178 snake pipefish from a large portion of their range. Cytochrome *b* yielded 94 unique haplotypes ($H_d = 0.967 \pm 0.006$ S.D.) and 82 polymorphic sites ($\pi = 0.0050 \pm 0.0002$ S.D., Table 2).

Sequence analyses of 385 bp of the control region locus in 160 snake pipefish revealed variation in a dinucleotide microsatellite repeat in the control region at site 287.

Most sequences contained [TA]₁₀ but [TA]₉ occurred at a low frequency (0.16) and was present in all geographical regions analysed except the Norwegian Sea samples that failed to amplify at this locus (Table 1). Inclusion of a mitochondrial microsatellite is problematic for several reasons (Lunt, Whipple & Hyman, 1998). For example, microsatellites generally have much higher mutation rates (in terms of the gain or loss of a repeat) than nucleotide substitutions, and mutation rates in general are unknown for mitochondrial microsatellites (Sia, Butler, Dominska *et al.*, 2000) and for this microsatellite in particular. Moreover, we did not determine if heteroplasmy, or the coexistence of nonidentical mtDNA molecules in the same individual, is occurring in this species. Heteroplasmy is common in species that have microsatellites in mtDNA in the AT-rich or control region causing difficulties for population genetics analyses (Lunt *et al.*, 1998; Mayer & Kerth, 2005). Additionally, length variation in the microsatellite can be considered a gap or missing data and thus may be inappropriate for some analyses (Yang & Rannala, 2012). Therefore, we excluded the variable repeat from all analyses, and we resolved 383 bps of the control region resulting in 63 unique haplotypes ($H_d = 0.893 \pm 0.015$ S.D.) and 43 polymorphic sites ($\pi = 0.0054 \pm 0.0003$ S.D., Table 2).

Population structure

A maximum likelihood haplotype network constructed for cytochrome *b* showed four major ($n \geq 10$) haplotypes (Fig. 2a). However, these haplotypes were comprised of representative individuals from most regions such that no genetic clustering could be discerned. For the control region, a maximum likelihood haplotype network also showed four major haplotypes and individuals from all populations were represented in these major haplotypes (Fig. 3). Overall, haplotype networks showed a star-like topology with high numbers of low-frequency mutations representative of a recent population expansion.

Pairwise ϕ_{ST} values ranged from -0.119 to 0.212 with a mean of -0.009 for cytochrome *b*, and ranged from -0.290 to 0.242 with a mean of -0.030 for the control region (Appendix S1). No significant differences between pairs of collections were detected after Bonferroni adjustment to correct for multiple comparisons (Rice, 1989). An AMOVA among collections within regions pooled across years indicated no significant population structuring (ϕ_{ST}) for either cytochrome *b* or the control region (Table 3, Appendix S1). Finally, no evidence for genetically distinct ecotypes or cryptic species was found when investigating individuals captured in nearshore benthic habitats versus pelagic captures (Table 3). In all AMOVA comparisons, the variance explained by among groups and among collections within groups was negligible in comparison to the variance within collections (Table 3).

Population expansion

Significant values of Fu's F_s , Tajima's D , and Ramos-Onsins and Rozas's R_2 found in cytochrome *b* sequences, with the exception of the Spanish coast collection, support a scenario of recent population expansion when samples were pooled within locations and sampling times (Table 2). The control region, on the other hand, showed significant F_s values, but only Continental shelf and French Coast collections showed significant departures from neutrality in Tajima's D and R_2 tests (Table 2). When collections were pooled across all location and sampling times, both the cytochrome *b* and control region showed highly significant F_s , D and R_2 values indicating recent population expansion was apparent (Table 2). Mismatch distributions were unimodal and failed to reject the hypothesis of the sudden expansion model (Appendix S1).

Bayesian Skyline Plot analysis with concatenated cytochrome *b* and control region sequences revealed that snake pipefish experienced a historical effective population size expansion starting about 100ka and obtaining current effective population sizes around 50ka

(Fig. 3). Individual effective population size analyses showed expansions approximately 125ka for cytochrome *b* and 40ka for control region sequences (Appendix S1).

Discussion

Our study provides insight into the phylogeographic history of the snake pipefish, a widely distributed syngnathid species in the northeastern Atlantic with a pelagic lifestyle. Results from molecular analyses did not reveal any clear patterns in population structure by regions, despite relatively high haplotype diversity estimates. However, we did uncover a signature of a Pleistocene population expansion in the northeastern Atlantic approximately 50-100ka. We also did not find any evidence for genetically distinct coastal or pelagic ecotypes or cryptic species, indicating that differences in phenotype are likely due to differences in ecological conditions. Taken together, our results point to a large population of snake pipefish throughout its contemporary northeastern Atlantic distribution and that such a high degree of homogenization is likely the result of a combination of specialized/unique life history traits and mixing by oceanic currents.

Historical phylogeography

The history of the northeastern Atlantic is one of extreme climatic changes, with multiple glaciation cycles until the late Pleistocene (last interglacial period ~125 ka, last glacial maximum ~20ka, Mokeddem, McManus & Oppo, 2014). During this period, many marine species were deeply impacted by glacial activity, which caused large reductions and/or shifts in suitable habitat (Mäkinen & Merilä, 2008) occasionally leading to precipitous population declines (Almada, Pereira, Robalo *et al.*, 2008; Boehme, Thompson, Fedak *et al.*, 2012). The subsequent recolonization of the northeastern Atlantic oft times leads to complex genetic signatures of glacial refugia, range expansions and bottlenecks (e.g. Coyer, Peters, Stam *et al.*, 2003; Gysels, Hellemans, Pampoulie *et al.*, 2004; Luttikhuizen, Campos, van

303 Bleijswijk *et al.*, 2008; Maggs, Castilho, Foltz *et al.*, 2008; Robalo, Castilho, Francisco *et al.*,
 304 2012).

305 The current study suggests that the snake pipefish underwent a population expansion
 306 in the northeastern Atlantic Ocean approximately 50-100ka during the Pleistocene. This
 307 scenario is supported by a star-like network topology, mismatch distribution analyses and an
 308 increase in effective population size indicating a recent expansion and/or a short evolutionary
 309 history of the species in the northeastern Atlantic Ocean (Grant & Bowen, 1998). Several
 310 other species show limited geographic partitioning and a sudden population expansion much
 311 earlier than the last glacial maximum similar to the snake pipefish. For example, the time of
 312 population expansion in the snake pipefish as estimated by BSP is similar to that for Atlantic
 313 Cod (*Gadus morhua*) which shows population expansion ~60ka in the northeastern Atlantic
 314 based on mismatch distributions of the mitogenome (Carr & Marshall, 2008). Other species
 315 that show signatures of population expansion predating the last glacial maximum include
 316 pelagic migrating species such as Atlantic bluefin tuna *Thunnus thynnus* (Bremer, Viñas,
 317 Mejuto *et al.*, 2005) and the European anchovy, *Engraulis encrasicolus* (Silva, Horne &
 318 Castilho, 2014). In contrast, other sympatric nearshore species of syngnathids closely
 319 associated with seagrass habitats show more recent recolonization of the northeastern
 320 Atlantic (15-36ka) based on coalescence analysis (Wilson & Eigenmann Veraguth, 2010).
 321 Thus, it appears that pelagic species show a reduced influence of Pleistocene glacial cycles in
 322 the northeastern Atlantic.

323 Our estimates of expansion time are likely conservative, given that mutation rates are
 324 time dependent and are higher in younger lineages (Ho, Phillips, Cooper *et al.*, 2005; Ho,
 325 Lanfear, Bromham *et al.*, 2011; Crandall, Sbrocco, DeBoer *et al.*, 2012; Grant, 2015). It is
 326 therefore possible that time since expansion is more recent than estimated from BSP. It is also
 327 probable that the differences in mutation rates chosen for cytochrome *b* and the control region

contribute to the discrepancy between individual locus estimates of expansion time and may have led to a more gradual expansion time using the concatenated loci as compared to individual loci (Appendix S1). Fluctuations in drift, in population size, selection, and reproductive skew may also influence divergence times (Burridge, Craw, Fletcher *et al.*, 2008; Grant, 2015; Niwa, Nashida & Yanagimoto, 2016). Finally, results of the BSP do not show recent (i.e., 10ka) fluctuations in effective population size although the probability of mtDNA markers capturing these phenomena is extremely unlikely.

Population structure

Population structuring over a large geographical range was not evident in this species from ϕ_{ST} and AMOVA results. These results mirror those reported for several Atlantic fish species where high mobility and/or high dispersal potential are highlighted as causes for little or no population sub-structuring (Nesbo, Rueness, Iversen *et al.*, 2000; Dannewitz, Maes, Johansson *et al.*, 2005; Carr & Marshall, 2008; Limborg, Hanel, Debes *et al.*, 2012). Our results suggest that despite having relatively low swimming ability, the snake pipefish is capable of dispersing over long distances, likely aided by a pelagic lifestyle and faster development rates relative to other northeastern Atlantic syngnathid species (Braga Goncalves, Ahnesjö & Kvarnemo, 2016). A similar pattern has been described in the non-migratory, demersal marine fish, *Sebastes schegellii*, where the association of larvae and juveniles with rafting precludes population genetic differentiation throughout its geographical range (Zhang, Yanagimoto, Zhang *et al.*, 2016). These results provide a stark contrast with several species of syngnathid that show population substructure over large spatial scales using mitochondrial markers (Lourie & Vincent, 2004; Lourie *et al.*, 2005; Teske *et al.*, 2005; Mobley *et al.*, 2010; Wilson & Eigenmann Veraguth, 2010; Wilson *et al.*, 2016).

Greater resolution of population sub-structuring in snake pipefish could potentially be provided by additional nuclear markers. However, attempts to design microsatellite markers

for this species have not yet yielded sufficient numbers of polymorphic microsatellites to conduct such analyses (KB Mobley, I Braga Goncalves, unpublished data). Future studies that benefit from combined mtDNA and nuclear markers may reveal additional insights into the phylogeography of this species including resolution of the time since population expansion and estimate gene flow within the species.

Range expansion

Although a significant number of snake pipefish were caught in polar regions during the population increase in 2003-2007, they may have originated in the southern Norwegian Sea and drifted to these locations in ocean currents (Nesbo *et al.*, 2000; Luttikhuizen *et al.*, 2008). In our study, we obtained only a few samples in polar waters from the Norwegian Sea in 2008. Haplotypes obtained from these samples were not distinct from southern populations suggesting that they are derived from the same large Atlantic gene pool. Due to the low abundance of snake pipefish after the population increase in north Atlantic waters (Heath *et al.*, 2012), future surveys should investigate whether or not these fish continue to inhabit polar regions representing a true range expansion or whether these were just transient individuals that rafted northward on ocean currents during the 2003-2007 outbreak.

Cryptic species

Previous studies have documented phenotypic differences between snake pipefish collected in coastal and oceanic areas (Holt & Byrne, 1906; Zhang *et al.*, 2016). Based on these differences, two species have been proposed previously, *E. aequoreus* found in oceanic waters and *E. anguineus* found in inshore habitats (Yarrel, 1839; Moreau, 1881), although *E. anguineus* is not currently recognized as a valid species (Dawson, 1985). These oceanic and coastal ‘ecotypes’ were recently described and a third potential intermediate ecotype that shares phenotypic similarities with both coastal and oceanic forms has been proposed (van Damme & Couperus, 2008). Despite differences in coloration and body condition, van

Damme and Couperus (2008) found no differences in ring and fin ray counts between perceived ecotypes in either sex and therefore concluded that all specimens belong to a single species. Based on mitochondrial haplotype frequencies, we also find no support that oceanic pelagic and benthic coastal ecotypes of snake pipefish collected in these habitats are different species. Instead, the snake pipefish appears to form a single species in the northeastern Atlantic and phenotypic plasticity in response to local ecological conditions encountered is the most probable explanation for the presence of multiple ecotypes.

Conservation concerns

According to the International Union for Conservation and Nature red list, *E. aequoreus* is evaluated as least concern (IUCN, 2017). Despite this listing, the sudden and substantial population increase in snake pipefish, although short lived, had a dramatic effect on the ecology of the northeastern Atlantic during the outbreak, and therefore warrants special mention. The significant increase in the numbers of snake pipefish in European waters was paralleled by a similar increase in the number of pipefish fed by parents to seabird nestlings in several species around the UK, Norway, Iceland, and the Faroe Islands (Luttikhuizen *et al.*, 2008; Anderson, Evans, Potts *et al.*, 2014). The hard exoskeleton and relatively low nutritional content of the snake pipefish (Harris, Newell, Daunt *et al.*, 2008) make them unsuitable for consumption by nestlings and adults alike, and their increased use as food items was associated with seabird breeding failures in the UK (Mavor, Parsons, Heubeck *et al.*, 2005; Mavor, Parsons, Heubeck *et al.*, 2006; Luttikhuizen *et al.*, 2008).

Despite the ecological significance of this species, the cause of the recent population increase of the snake pipefish still remains unknown, as does its subsequent decline (Heath *et al.*, 2012). Several non-mutually exclusive hypotheses have been put forward to explain its

population increase and expansion, namely: 1) a rise in surface seawater temperatures promoting longer breeding seasons and higher recruitment (Kirby *et al.*, 2006; Gremillet & Boulinier, 2009; Neumann, Ehrich & Kroncke, 2009; Anderson *et al.*, 2014), 2) a by-product of tracking changing and/or shifting plankton communities (van Damme & Couperus, 2008), 3) promoted by the establishment of invasive algal species such as the Japanese seaweed, *Sargassum muticum*, that increased the amount of suitable habitat for successful reproduction in the coastal regions (Gysels *et al.*, 2004) and, 4) a result of decreasing population size of interspecific competitors such as Lesser Sandeels (*Ammodytes marinus*) due to fishing and climate change (Heath *et al.*, 2012; Anderson *et al.*, 2014). Yet, there is no conclusive evidence to explain the sudden expansion.

The increase in snake pipefish potentially started in one source spot on the continental shelf and dispersed over the entire northeastern Atlantic. Interestingly, the increase and expansion in the pelagic environment itself can be seen as a strategy of the species to colonize new suitable habitats. However, expansions of the same magnitude appear to be very rare: there are indications that a similar population increase took place at the end of the nineteenth century (Brongersma-Sanders, 1957; van Damme & Couperus, 2008). Climate change is expected to change ocean currents which, together with higher sea surface temperatures, will affect larval import, export and recruitment, leading to faster development, shorter larval stages and dispersal into new habitats (Kendall, Poti & Karnauskas, 2016). Improving our understanding of snake pipefish reproductive biology, habitat, feeding ecology, and dispersal potential is a critical next step to help pinpoint how current and future changes in climate and in prey distributions in the northeastern Atlantic may potentiate further population increases, which in turn may affect community structure.

Conclusions

Understanding species' current distribution patterns and historical demography is a fundamental goal in evolutionary biology. Our study contributes to this goal by investigating phylogeographic patterns in a species that has undergone a sudden population increase and range expansion that had a negative impact on the ecosystem of the northeastern Atlantic. Although the cause for the recent population increase and range expansion and contraction are still unknown, the phylogeographic patterns uncovered in our study demonstrate that the snake pipefish represents a single large population with no evidence of population substructuring. This result, in contrast to all other syngnathids studied to date, may be explained by the pelagic lifestyle and poor swimming capabilities of the species, allowing individuals to be transported long distances by ocean currents. Our study adds to the understanding of this ecologically important species and future studies should incorporate a wider range of genetic markers to investigate population demographics, particularly concerning the recent population increase.

Supplementary Material

Additional Supporting Information may be found in the online version of this article:
Appendix S1: Population pairwise Φ_{ST} estimates, mismatch distributions, and individual Bayesian skyline plots for cytochrome *b* and control region loci.

Acknowledgements

We would like to thank Anders Berglund, Zach Cress, Loren Mendoza, Laurent Leveque, Ian Probert and the staff at the Roscoff Marine Station Centre, France and for assistance with field collections. We would also like to thank Adam Jones and Clay Small at Texas A&M University, Tammes Menne at the Zoological Museum, University of Copenhagen and Tony Wilson and Alexandra Wegmann at the University of Zürich for access to preserved

specimens. Sophie Bodenstein, Thoomka Brüning, Withe Derner, Nadja Gogrefe, Sybille Liedtke, and Ines Schlultz assisted with DNA sequencing. We thank Folmer Bokma at Umeå University and the Max Planck Society for technical support. We thank two anonymous reviewers for helpful comments on the manuscript. This work was supported by a transnational access grant to KM and IBG from the Association of European Marine Biological Laboratories (ASSEMBLE), for fieldwork at Roscoff Marine Station, France and Lundgrenska Stiftelserna and Helge Axelsson Johnsons stiftelse for financial support to IBG.

Data Accessibility

All cytochrome *b* and control region sequences will be available in Genbank and supporting sample information and haplotype lists will be deposited in Dryad.

References

- Almada VC, Pereira AM, Robalo JI, Fonseca JP, Levy A, Maia C, Valente A. 2008.** Mitochondrial DNA fails to reveal genetic structure in sea-lampreys along European shores. *Molecular Phylogenetics and Evolution* **46**: 391-396.
- Anderson HB, Evans PGH, Potts JM, Harris MP, Wanless S. 2014.** The diet of Common Guillemot *Uria aalge* chicks provides evidence of changing prey communities in the North Sea. *Ibis* **156**: 23-34.
- Avice JC. 2000.** *Phylogeography: the History and Formation of Species*. Harvard University Press: Cambridge, MA.
- Bernard AM, Feldheim KA, Heithaus MR, Wintner SP, Wetherbee BM, Shivji MS. 2016.** Global population genetic dynamics of a highly migratory, apex predator shark. *Molecular Ecology* **25**: 5312-5329.

- 475 **Boehme L, Thompson D, Fedak M, Bowen D, Hammill MO, Stenson GB. 2012.** How
 476 many seals were there? The global shelf loss during the Last Glacial Maximum and its
 477 effect on the size and distribution of grey seal populations. *PLoS One* **7**: e53000.
- 478 **Bowen BW, Muss A, Rocha LA, Grant WS. 2006.** Shallow mtDNA coalescence in Atlantic
 479 pygmy angelfishes (genus *Centropyge*) indicates a recent invasion from the Indian
 480 Ocean. *Journal of Heredity* **96**: 1-12.
- 481 **Braga Goncalves I, Ahnesjö I, Kvarnemo C. 2016.** Evolutionary ecology of pipefish
 482 brooding structures: embryo survival and growth do not improve with a pouch.
 483 *Ecology and Evolution* **6**: 6308-3620.
- 484 **Bremer JRA, Viñas J, Mejuto J, Ely B, Pla C. 2005.** Comparative phylogeography of
 485 Atlantic bluefin tuna and swordfish: the combined effects of vicariance, secondary
 486 contact, introgression, and population expansion on the regional phylogenies of two
 487 highly migratory pelagic species. *Molecular Phylogenetics and Evolution* **36**: 169-
 488 187.
- 489 **Brongersma-Sanders M. 1957.** Mass mortality in the sea. In: Hedgpeth JW, ed. *Treatise on*
 490 *Marine Ecology and Paleoecology I*. Geological Society of America Memoir. 941-
 491 1010.
- 492 **Burridge CP, Craw D, Fletcher D, Waters JM. 2008.** Geological dates and molecular
 493 rates: fish DNA sheds light on time dependency. *Molecular Biology and Evolution*
 494 **25**: 624-633.
- 495 **Carr SM, Marshall HD. 2008.** Intraspecific phylogeographic genomics from multiple
 496 complete mtDNA Genomes in Atlantic cod (*Gadus morhua*): Origins of the

- 497 "Codmother," transatlantic vicariance and midglacial population expansion. *Genetics*
 498 **180**: 381-389.
- 499 **Chenoweth SF, Hughes JM, Connolly RM. 2002.** Phylogeography of the pipefish,
 500 *Urocampus carinirostris*, suggests secondary intergradation of ancient lineages.
 501 *Marine Biology* **141**: 541-547.
- 502 **Coyer JA, Peters AF, Stam WT, Olsen JL. 2003.** Post-ice age recolonization and
 503 differentiation of *Fucus serratus* L. (Phaeophyceae; Fucaceae) populations in
 504 Northern Europe. *Molecular Ecology* **12**: 1817-1829.
- 505 **Crandall ED, Sbrocco EJ, DeBoer TS, Barber PH, Carpenter KE. 2012.** Expansion
 506 dating: calibrating molecular clocks in marine species from expansions onto the
 507 Sunda shelf following the Last Glacial Maximum. *Molecular Biology and Evolution*
 508 **29**: 707-719.
- 509 **Dannewitz J, Maes GE, Johansson L, Wickstrom H, Volckaert FAM, Jarvi T. 2005.**
 510 Panmixia in the European eel: a matter of time. *Proceedings of the Royal Society B-*
 511 *Biological Sciences* **272**: 1129-1137.
- 512 **Darriba D, Taboada GL, Doallo R, Posada D. 2012.** jModelTest 2: more models, new
 513 heuristics and parallel computing. *Nature Methods* **9**: 772-772.
- 514 **Dawson CE. 1985.** *Indo-Pacific pipefishes*. Gulf Coast Research Lab: Ocean Springs.
- 515 **Dawson CE. 1986.** Syngnathidae. In: Whitehead PJP, Bauchot M-L, Hureau J-C, Nielsen J
 516 and Tortonese E, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*.
 517 UNESCO, Paris. 628-639.

- 518 **Drummond AJ, Rambaut A. 2007.** BEAST: Bayesian evolutionary analysis by sampling
519 trees. *BMC Evolutionary Biology* **7**: 1-8.
- 520 **Drummond AJ, Rambaut A, Shapiro B, Pybus OG. 2005.** Bayesian coalescent inference
521 of past population dynamics from molecular sequences. *Molecular Biology and*
522 *Evolution* **22**: 1185-1192.
- 523 **Dunker G. 1915.** Revision fer Sungnathidae. *Mitteilungen aus dem Naturhistorischen*
524 *Museum in Hamburg* **32**: 32-34.
- 525 **Edgar RC. 2004.** MUSCLE: multiple sequence alignment with high accuracy and high
526 throughput. *Nucleic Acids Research* **32**: 1792-1797.
- 527 **Excoffier L, Laval G, Schneider S. 2005.** Arelquin (version 3.0): an integrated software
528 package for population genetics data analysis. *Evolutionary Bioinformatics Online* **1**:
529 47-50.
- 530 **Excoffier L, Smouse PE, Quattro JM. 1992.** Analysis of molecular variance inferred from
531 metric distances among DNA haplotypes: application to human mitochondrial DNA
532 restriction data. *Genetics* **131**: 479-491.
- 533 **Fedrizzi N, Stiassny MLJ, Boehm JT, Dougherty ER, Amato G, Mendez M. 2015.**
534 Population genetic structure of the dwarf seahorse (*Hippocampus zosterae*) in Florida.
535 *PLoS One* **10**: 1-13.
- 536 **Felsenstein J. 1989.** PHYLIP – Phylogeny Inference Package (version 3.2). *Cladistics* **5**:
537 164-166.

- 538 **Fleischer D, Schaber M, Piepenburg D. 2007.** Atlantic snake pipefish (*Entelurus*
539 *aequoreus*) extends its northward range to Svalbard (Arctic Ocean). *Polar Biology* **30**:
540 1359-1362.
- 541 **Fries B, Ekström CU, Sundevall C. 1895.** *A history of Scandinavian fishes*. Norstedt &
542 Söner: Stockholm.
- 543 **Fu YX. 1997.** Statistical tests of neutrality of mutations against population growth,
544 hitchhiking and background selection. *Genetics* **147**: 915-925.
- 545 **Grant WS. 2015.** Problems and cautions with sequence mismatch analysis and Bayesian
546 skyline plots to infer historical demography. *Journal of Heredity* **106**: 333-346.
- 547 **Grant WS, Bowen BW. 1998.** Shallow population histories in deep evolutionary lineages of
548 marine fishes: insights from sardines and anchovies and lessons for conservation.
549 *Journal of Heredity* **89**: 415-426.
- 550 **Grantham BA, Eckert GL, Shanks AL. 2003.** Dispersal potential of marine invertebrates in
551 diverse habitats. *Ecological Applications* **13**: S108-S116.
- 552 **Gremillet D, Boulinier T. 2009.** Spatial ecology and conservation of seabirds facing global
553 climate change: a review. *Marine Ecology Progress Series* **391**: 121-137.
- 554 **Gysels ES, Hellemans B, Pampoulie C, Volckaert FAM. 2004.** Phylogeography of the
555 common goby, *Pomatoschistus microps*, with particular emphasis on the colonization
556 of the Mediterranean and the North Sea. *Molecular Ecology* **13**: 403-417.
- 557 **Harris MP, Beare D, Toresen R, Nottestad L, Kloppmann M, Dorner H, Peach K,**
558 **Rushton DRA, Foster-Smith J, Wanless S. 2007.** A major increase in snake pipefish

- 559 (*Entelurus aequoreus*) in northern European seas since 2003: potential implications
560 for seabird breeding success. *Marine Biology* **151**: 973-983.
- 561 **Harris MP, Newell M, Daunt F, Speakman JR, Wanless S. 2008.** Snake Pipefish
562 *Entelurus aequoreus* are poor food for seabirds. *Ibis* **150**: 413-415.
- 563 **Hasse JJ. 1974.** Observations on egg hatching of *Syngnathoides biaculeatus* (Bloch).
564 *Micronesia* **10**: 279-283.
- 565 **Heath MR, Neat FC, Pinnegar JK, Reid DG, Sims DW, Wright PJ. 2012.** Review of
566 climate change impacts on marine fish and shellfish around the UK and Ireland.
567 *Aquatic Conservation-Marine and Freshwater Ecosystems* **22**: 337-367.
- 568 **Henderson PA, Bird DJ. 2010.** Fish and macro-crustacean communities and their dynamics
569 in the Severn Estuary. *Marine Pollution Bulletin* **61**: 100-114.
- 570 **Hewitt GM. 1999.** Post-glacial re-colonization of European biota. *Biological Journal of the*
571 *Linnean Society* **68**: 87-112.
- 572 **Ho SYW, Lanfear R, Bromham L, Phillips MJ, Soubrier J, Rodrigo G, Cooper A. 2011.**
573 Time-dependent rates of molecular evolution. *Molecular Ecology* **24**: 6007-6012.
- 574 **Ho SYW, Phillips MJ, Cooper A, Drummond AJ. 2005.** Time dependency of molecular
575 rate estimates and systematic overestimation of recent divergence times. *Molecular*
576 *Biology and Evolution* **22**: 1561-1568.
- 577 **Holt EWL, Byrne LW. 1906.** First report on the fishes of the Irish Atlantic slope. *Fisheries*
578 *Ireland Scientific Investigations* **II**: 1-28.

- 579 **IUCN. 2017.** The IUCN Red List of Threatened Species. Version 2017-1.
 580 www.iucnredlist.org. Downloaded on 15 August 2017.
- 581 **Kendall MS, Poti M, Karnauskas KB. 2016.** Climate change and larval transport in the
 582 ocean: fractional effects from physical and physiological factors. *Global Change*
 583 *Biology* **22**: 1532-1547.
- 584 **Kirby RR, Johns DG, Lindley JA. 2006.** Fathers in hot water: rising sea temperatures and a
 585 Northeastern Atlantic pipefish baby boom. *Biology Letters* **2**: 597-600.
- 586 **Kloppmann MHF, Ulleweit J. 2007.** Off-shelf distribution of pelagic snake pipefish,
 587 *Entelurus aequoreus* (Linnaeus, 1758), west of the British Isles. *Marine Biology* **151**:
 588 271-275.
- 589 **Kocher TD, K. TW, Meyer A, Edwards SV, Pääbo S, Villablan CA, Villablan FX,**
 590 **Wilson AC. 1989.** Dynamics of mitochondrial DNA evolution in animals:
 591 amplification and sequencing with conserved primers. *Proceedings of the National*
 592 *Academy of Sciences USA* **74**: 5436-5467.
- 593 **Librado P, Rozas J. 2009.** A software for comprehensive analysis of DNA polymorphism
 594 data. *Bioinformatics* **25**: 1451-1452.
- 595 **Limborg MT, Hanel R, Debes PV, Ring AK, Andre C, Tsigenopoulos CS, Bekkevold D.**
 596 **2012.** Imprints from genetic drift and mutation imply relative divergence times across
 597 marine transition zones in a pan-European small pelagic fish (*Sprattus sprattus*).
 598 *Heredity* **109**: 96-107.

- 599 **Lourie SA, Green DM, Vincent ACJ. 2005.** Dispersal, habitat differences, and comparative
600 phylogeography of Southeast Asian seahorses (Syngnathidae: *Hippocampus*).
601 *Molecular Ecology* **14**: 1073-1094.
- 602 **Lourie SA, Vincent ACJ. 2004.** A marine fish follows Wallace's Line: the phylogeography
603 of the three-spot seahorse (*Hippocampus trimaculatus*, Syngnathidae, Teleostei) in
604 Southeast Asia. *Journal of Biogeography* **31**: 1975-1985.
- 605 **Lunt DH, Whipple LE, Hyman BC. 1998.** Mitochondrial DNA variable number tandem
606 repeats (VNTRs): utility and problems in molecular ecology. *Molecular Ecology* **7**:
607 1441-1455.
- 608 **Luttikhuizen PC, Campos J, van Bleijswijk J, Peijnenburg K, van der Veer HW. 2008.**
609 Phylogeography of the common shrimp, *Crangon crangon* (L.) across its distribution
610 range. *Molecular Phylogenetics and Evolution* **46**: 1015-1030.
- 611 **Maggs CA, Castilho R, Foltz D, Henzler C, Jolly MT, Kelly J, Olsen J, Perez KE, Stam
612 W, Vainola R, Viard F, Wares J. 2008.** Evaluating signatures of glacial refugia for
613 north Atlantic benthic marine taxa. *Ecology* **89**: S108-S122.
- 614 **Mäkinen HS, Merilä J. 2008.** Mitochondrial DNA phylogeography of the three-spined
615 stickleback (*Gasterosteus aculeatus*) in Europe - Evidence for multiple glacial refugia.
616 *Molecular Biology and Evolution* **46**: 167-182.
- 617 **Mavor RA, Parsons M, Heubeck M, Schmitt S. 2005.** *Seabird numbers and breeding
618 success in Britain and Ireland, 2004*. Peterborough: Joint Nature Conservation
619 Committee (UK Nature Conservation).

- 620 **Mavor RA, Parsons M, Heubeck M, Schmitt S. 2006.** *Seabird numbers and breeding*
 621 *success in Britain and Ireland, 2005.* Peterborough: Joint Nature Conservation
 622 Committee (UK Nature Conservation).
- 623 **Mayer F, Kerth G. 2005.** Microsatellite evolution in the mitochondrial genome of
 624 Bechstein's bat (*Myotis bechsteinii*). *Journal of Molecular Evolution* **61**: 408-416.
- 625 **Meyer A, Kocher TD, Basasibwaki P, Wilson AC. 1990.** Monophyletic origin of Lake
 626 Victoria cichlid fishes suggested by mitochondrial DNA sequences. *Nature* **347**: 550-
 627 553.
- 628 **Mi PT, Kornienko Es, Drozdov AL. 1998.** Embryonic and larval development of the
 629 seahorse *Hyppocampus kuda*. *Russian Journal of Marine Biology* **24**: 325-329.
- 630 **Mobley KB, Small CM, Jones AG. 2011.** The genetics and genomics of Syngnathidae
 631 (pipefishes, seahorses and seadragons). *Journal of Fish Biology* **78**: 1624-1646.
- 632 **Mobley KB, Small CM, Jue NK, Jones AG. 2010.** Population structure of the dusky
 633 pipefish (*Syngnathus floridae*) from the Atlantic and Gulf of Mexico, as revealed by
 634 mitochondrial DNA and microsatellite analyses. *Journal of Biogeography* **37**: 1363-
 635 1377.
- 636 **Mokeddem Z, McManus JF, Oppo DW. 2014.** Oceanographic dynamics and the end of the
 637 last interglacial in the subpolar North Atlantic. *Proceedings of the National Academy*
 638 *of Sciences of the U.S.A.* **111**: 11263-11268.
- 639 **Monteiro NM, Almada VC, Vieira MN. 2003.** Early life history of the pipefish *Nerophis*
 640 *lumbriciformis* (Pisces : Syngnathidae). *Journal of the Marine Biological Association*
 641 *of the United Kingdom* **83**: 1179-1182.

- 642 **Monteiro NM, Berglund A, Vieira MN, Almada VC. 2006.** Reproductive migrations of the
 643 sex role reversed pipefish *Nerophis lumbriciformis* (Pisces; Syngnathidae). *Journal of*
 644 *Fish Biology* **69**: 66-74.
- 645 **Moran EV, Alexander JM. 2014.** Evolutionary responses to global change: lessons from
 646 invasive species. *Ecology Letters* **17**: 637-649.
- 647 **Moreau E. 1881.** *Histoire Naturelle des Poissons de la France*. Libraire de l' academie de
 648 Medecine: Paris.
- 649 **Nei M. 1987.** *Molecular evolutionary genetics*. Columbia University Press: New York.
- 650 **Nesbo CL, Rueness EK, Iversen SA, Skagen DW, Jakobsen KS. 2000.** Phylogeography
 651 and population history of Atlantic mackerel (*Scomber scombrus* L.): a genealogical
 652 approach reveals genetic structuring among the eastern Atlantic stocks. *Proceedings*
 653 *of the Royal Society B-Biological Sciences* **267**: 281-292.
- 654 **Neumann H, Ehrich S, Kroncke I. 2009.** Variability of epifauna and temperature in the
 655 northern North Sea. *Marine Biology* **156**: 1817-1826.
- 656 **Nickel J, Cursons R. 2012.** Genetic diversity and population structure of the pot-belly
 657 seahorse *Hippocampus abdominalis* in New Zealand. *New Zealand Journal of Marine*
 658 *and Freshwater Research* **46**: 207-218.
- 659 **Niwa H-S, Nashida K, Yanagimoto T. 2016.** Reproductive skew in Japanese sardine
 660 inferred from DNA sequences. *ICES Journal of Marine Science* **73**: 2181-2189.
- 661 **Olsen JL, Stam WT, Coyer JA, Reusch TBH, Billingham M, Bostrom C, Calvert E,**
 662 **Christie H, Granger S, La Lumiere R, Milchakova N, Oudot-Le Secq MP,**
 663 **Procaccini G, Sanjabi B, Serrao E, Veldsink J, Widdicombe S, Wyllie-**

- 664 **Echeverria S. 2004.** North Atlantic phylogeography and large-scale population
665 differentiation of the seagrass *Zostera marina* L. *Molecular Ecology* **13**: 1923-1941.
- 666 **OSPAR. 2010.** *Quality Status Report 2010*. OSPAR Commission: London.
- 667 **Pääbo S, Thomas WK, Whitfield KM, Kumazawa Y, Wilson AC. 1991.** Rearrangements
668 of mitochondrial transfer RNA genes in marsupials. *Journal of Molecular Evolution*
669 **33**: 426-430.
- 670 **Planas M, Blanco A, Chamorro A, Valladares S, Pintado J. 2012.** Temperature-induced
671 changes of growth and survival in the early development of the seahorse
672 *Hippocampus guttulatus*. *Journal of Experimental Marine Biology and Ecology* **438**:
673 154-162.
- 674 **Polte P, Buschbaum C. 2008.** Native pipefish *Entelurus aequoreus* are promoted by the
675 introduced seaweed *Sargassum muticum* in the northern Wadden Sea, North Sea.
676 *Aquatic Biology* **3**: 11-18.
- 677 **Rambaut A, Suchard MA, Xie D, Drummond AJ. 2014.** Tracer v1.6, Available from
678 <http://beast.bio.ed.ac.uk/Tracer>.
- 679 **Ramírez-Soriano A, Ramos-Onsins SE, Rozas J, Calafell F, Navarro A. 2008.** Statistical
680 power analysis of neutrality tests under demographic expansions, contractions and
681 bottlenecks with recombination. *Genetics* **179**: 555-567.
- 682 **Ramos-Onsins SE, Rozas J. 2002.** Statistical properties of new neutrality tests against
683 population growth. *Molecular Biology and Evolution* **19**: 2092-2100.
- 684 **Rice WR. 1989.** Analyzing tables of statistical tests. *Evolution* **43**: 223-225.

- 685 **Robalo JI, Castilho R, Francisco SM, Almada F, Knutsen H, Jorde PE, Pereira AM,**
686 **Almada VC. 2012.** Northern refugia and recent expansion in the North Sea: the case
687 of the wrasse *Symphodus melops* (Linnaeus, 1758). *Ecology and Evolution* **2**: 153-
688 164.
- 689 **Rusyaev SM, Dolgov AV, Karamushko OV. 2007.** Captures of snake pipefish *Entelurus*
690 *aequoreus* in the Barents and Greenland seas. *Journal of Ichthyology* **47**: 544-546.
- 691 **Salzburger W, Ewing GB, von Haeseler A. 2011.** The performance of phylogenetic
692 algorithms in estimating haplotype genealogies with migration. *Molecular Ecology*
693 **20**: 1952-1963.
- 694 **Schmitt T. 2007.** Molecular biogeography of Europe: Pleistocene cycles and postglacial
695 trends. *Frontiers in Zoology* **4**: 11.
- 696 **Sia EA, Butler CA, Dominska M, Greenwell P, Fox TD, Petes TD. 2000.** Analysis of
697 microsatellite mutations in the mitochondrial DNA of *Saccharomyces cerevisiae*.
698 *Proceedings of the National Academy of Sciences of the U.S.A.* **97**: 250-255.
- 699 **Silva G, Horne JB, Castilho R. 2014.** Anchovies go north and west without losing diversity:
700 post-glacial range expansions in a small pelagic fish. *Journal of Biogeography* **41**:
701 1171-1182.
- 702 **Silva K, Monteiro NM, Almada VC, Vieira MN. 2006.** Development and early life history
703 behaviour of aquarium reared *Syngnathus acus* (Pisces: Syngnathidae). *Journal of*
704 *Marine Biology Association UK* **86**: 1469-1472.

- 705 **Silva K, Monteiro NM, Vieira MN, Almada VC. 2006.** Reproductive behavior of the
 706 black-striped pipefish, *Syngnathus abaster* (Pisces; Syngnathidae). *Journal of Fish*
 707 *Biology* **69**: 1860-1869.
- 708 **Stiller J, Wilson NG, Donnellan S, Rouse GW. 2017.** The leafy seadragon, *Phycodurus*
 709 *eques*, a flagship species with low but structured genetic variability. *Journal of*
 710 *Heredity* **108**: 152-162.
- 711 **Tajima F. 1989.** Statistical method for testing the neutral mutation hypothesis by DNA
 712 polymorphism. *Genetics* **123**: 585-595.
- 713 **Teske PR, Hamilton H, Palsbøll PJ, Choo CK, Gabr H, Lourie SA, Santos M, Sreepada**
 714 **A, Cherry MI, Matthee CA. 2005.** Molecular evidence for long-distance
 715 colonization in an Indo-Pacific seahorse lineage. *Marine Ecology - Progress Series*
 716 **286**: 249-260.
- 717 **van Damme CJG, Couperus AS. 2008.** Mass occurrence of snake pipefish in the Northeast
 718 Atlantic: Result of change in climate? *Journal of Sea Research* **60**: 117-125.
- 719 **Vincent ACJ, Berglund A, Ahnesjö I. 1995.** Reproductive ecology of five pipefish species
 720 in one eelgrass meadow. *Environmental Biology of Fishes* **44**: 347-361.
- 721 **Wetzel JT, Wourms JP. 2004.** Embryogenesis in the dwarf seahorse *Hippocampus zosterae*
 722 (Syngnathidae). *Gulf and Caribbean Research* **16**: 27-35.
- 723 **Wilson AB, Eigenmann Veraguth I. 2010.** The impact of Pleistocene glaciation across the
 724 range of a widespread European coast species. *Molecular Ecology* **19**: 4535-4553.
- 725 **Wilson AB, Orr JW. 2011.** The evolutionary origins of Syngnathid fishes. *Journal of Fish*
 726 *Biology* **78**: 1603-1623.

- 727 **Wilson AB, Vincent A, Ahnesjö I, Meyer A. 2001.** Male pregnancy in seahorses and
 728 pipefishes (Family Syngnathidae): Rapid diversification of paternal brood pouch
 729 morphology inferred from a molecular phylogeny. *Journal of Heredity* **92**: 159-166.
- 730 **Wilson MJ, Vincent ACJ. 1998.** Preliminary success in closing the life cycle of exploited
 731 seahorse species, *Hippocampus* spp., in captivity. *Aquarium Sciences and*
 732 *Conservation* **2**: 179-196.
- 733 **Wilson NG, Stiller J, Rouse GW. 2016.** Barriers to gene flow in common seadragons
 734 (Syngnathidae: *Phyllopteryx taeniolatus*). *Conservation Genetics*: 1-14.
- 735 **Woodall LC, Koldewey HJ, Boehm JT, Shaw PW. 2015.** Past and present drivers of
 736 population structure in a small coastal fish, the European long snouted seahorse
 737 *Hippocampus guttulatus*. *Conservation Genetics* **16**: 1139-1153.
- 738 **Woodall LC, Koldewey HJ, Santos SV, Shaw PW. 2009.** First occurrence of the lined
 739 seahorse *Hippocampus erectus* in the eastern Atlantic Ocean. *Journal of Fish Biology*
 740 **75**: 1505-1512.
- 741 **Yang Z, Rannala B. 2012.** Molecular phylogenetics: principles and practice. *Nature Reviews*
 742 *Genetics* **13**: 303-314.
- 743 **Yarrel W. 1839.** Remarks on one species of the genus *Syngnathus*. *Annals of Natural*
 744 *History* **2**: 81-85.
- 745 **Zhang H, Yanagimoto T, Zhang XM, Song N, Gao TX. 2016.** Lack of population genetic
 746 differentiation of a marine ovoviviparous fish *Sebastes schlegelii* in Northwestern
 747 Pacific. *Mitochondrial DNA* **27**: 1748-1754.
- 748
- 749

750 Table 1. *Entelurus aequoreus* collections. Region of collection, number of individuals collected (n), year collected, location, collection
 751 abbreviation (AB), habitat, latitude, longitude, the number of individuals amplifiable at the cytochrome *b* locus (n_{cytb}) and control region locus
 752 (n_{CR}), and presence of the microsatellite repeat [TA]₉ in the control region are listed.

Region	n	Year	Location	AB	Habitat	Latitude	Longitude	n_{cytb}	n_{CR}	[TA] ₉
Continental Shelf	5	2005	Atlantic Ocean	CS1	pelagic	51°45'0.61"N	11°45'0.61"W	3	5	0
	2	2005	Atlantic Ocean	CS2	pelagic	56°10'2.39"N	9°47'4.81"W	2	1	0
	9	2005	Atlantic Ocean	CS3	pelagic	50°45'0.00"N	11°7'0.01"W	8	7	2
	15	2005	Atlantic Ocean	CS4	pelagic	51°15'1.80"N	13°42'1.80"W	14	13	3
	10	2007	Atlantic Ocean	CS5	pelagic	49°8'15.99"N	10°22'48"W	10	5	1
	23	2010	Atlantic Ocean	CS6	pelagic	51°52'42.60"N	13°8'3.01"W	22	21	3
Spanish Coast	10	2007	Galacia, ES	SC	coastal	42°15'N	8°52'W	10	8	1
North Sea	3	2005	North Sea	NS01	pelagic	57°48'34.99"N	0°52'50.70"W	3	1	0
	5	2005	North Sea	NS02	pelagic	57°52'7.14"N	3°14'54.06"W	5	2	2
	5	2005	North Sea	NS03	pelagic	57°44'55.03"N	1°21'28.26"W	5	2	0
	1	2005	North Sea	NS04	pelagic	58°10'10.99"N	0°33'42.23"W	1	1	0
	10	2005	North Sea	NS05	pelagic	58°10'37.27"N	3°10'12.00"W	10	7	5
	8	2007	North Sea	NS06	pelagic	56°21'38.99"N	2°4'58.19"W	8	8	1
	2	2007	North Sea	NS07	pelagic	56°7'34.21"N	3°28'3.00"E	2	2	0
	7	2007	North Sea	NS08	pelagic	53°28'53.40"N	0°54'54.00"E	7	7	1
	1	2007	North Sea	NS09	pelagic	55°36'16.20"N	2°46'31.19"E	1	1	0
	1	2007	North Sea	NS10	pelagic	55°53'26.41"N	4°15'45.61"E	1	1	0
	6	2007	North Sea	NS11	pelagic	56°45'29.41"N	1°33'30.60"W	6	5	0
	6	2007	North Sea	NS12	pelagic	55°23'28.79"N	1°34'34.79"E	6	6	0
Skagerrak/Kattegat	2	2003	Kølpn/Deget, DK	SK1	coastal	57°27'17.66"N	10°35'53.61"E	2	2	0
	1	2004	Dronningmølle, DK	SK2	coastal	56°6'6.07"N	12°24'34.77"E	1	1	0
	9	2005	Gåsö, SE	SK3	coastal	58°14'23.32"N	11°22'44.86"E	7	0	--

	35 ¹	2006	Gåsö, SE	SK4	coastal	58°14'23.32"N	11°22'44.86"E	20	19	2
Norwegian Sea	1	2008	Norwegian Sea	NOR1	pelagic	68°15'40.02"N	4° 7'45.00"E	0	0	--
	5	2008	Norwegian Sea	NOR2	pelagic	68°15'24.66"N	0°31'45.36"W	3	0	--
French Coast	55 ²	2010	Bay de Roscoff, FR	FC	coastal	48°42'13.75"N	3°55'11.67"W	21	35	4
Total	23									
	7							178	160	25

753 ¹A subset of 20 individuals was extracted. ²A subset of 35 individuals was extracted.

754 Table 2. Genetic diversity indices and tests of neutrality pooled across years in regions for Cytochrome *b* (900 bp) and control region (383 bp
 755 with microsatellite removed) loci. Number of individuals sequenced (n), number of unique haplotypes (H), haplotype diversity (H_d , \pm S.D.),
 756 nucleotide diversity (π , \pm S.D.), Tajima's *D* (*D*), Fu's *F_s* (*F_s*), and Ramos-Onsins and Rozas's *R₂* (*R₂*) test are given. Asterisks denote significant
 757 departures from neutrality: $P < 0.05 = *$, $P < 0.01 = **$, $P < 0.001 = ***$).

Region	Cytochrome <i>b</i>							Control region						
	n	H	H_d	π	<i>D</i>	<i>F_s</i>	<i>R₂</i>	n	H	H_d	π	<i>D</i>	<i>F_s</i>	<i>R₂</i>
Continental Shelf	59	41	0.972 (0.011)	0.0049 (0.0004)	-1.97**	-41.267***	0.038***	52	25	0.919 (0.022)	0.0052 (0.0004)	-2.04***	-76.50***	0.037***
Spanish Coast	10	7	0.911 (0.077)	0.0059 (0.0008)	-0.53	-0.89	0.127	8	6	0.929 (0.084)	0.0053 (0.0011)	0.25	-2.58*	0.023
North Sea	55	36	0.962 (0.015)	0.0048 (0.0004)	-1.82**	-31.99***	0.045***	43	24	0.942 (0.020)	0.0072 (0.0006)	-1.31	-19.64***	0.068
Skagarrak/Kattegat	30	22	0.968 (0.019)	0.0049 (0.0005)	-1.50*	-14.72***	0.063**	22	12	0.905 (0.044)	0.0064 (0.0009)	-1.27	-5.76***	0.079
Norwegian Sea	3	3	1.000 (0.272)	0.0030 (0.0009)	--	--	--	0	--	--	--	--	--	--
French Coast	21	19	0.986 (0.022)	0.0055 (0.0006)	-1.48*	-14.32***	0.069**	35	21	0.943 (0.024)	0.0074 (0.0009)	-1.70*	-16.11***	0.062*
Pooled	178	94	0.967 (0.006)	0.0050 (0.0002)	-2.22***	-133.75***	0.025***	160	63	0.893 (0.015)	0.0054 (0.0003)	-2.29***	-76.50***	0.025**

Table 3. Analysis of molecular variance (AMOVA) of *Entelurus aequoreus* based on 900 bp of mtDNA cytochrome *b* (Cytb) sequence and 383 bp of mtDNA control region (CR) sequence (microsatellite at position 287 excluded). The percent variation (% var), and P value are listed for among groups (Φ_{CT}), among collections within groups (Φ_{SC}) and within collections Φ_{ST} . AMOVAs were conducted with all collections within regions (regions) and all collections pooled within habitat (coastal vs. pelagic).

		Among groups			Among collections within groups			Within collections	
		% var	Φ_{CT}	P value	% var	Φ_{SC}	P value	Φ_{ST}	P value
All populations	Cytb	0.11	0.0011	0.4534	-0.62	-0.0062	0.5535	-0.0051	0.6189
	CR	3.76	0.0376	0.0603	-3.11	-0.0324	0.9151	0.0065	0.4495
Habitat	Cytb	-0.52	-0.0052	0.7534	-0.28	-0.0028	0.4951	-0.0080	0.6165
	CR	2.82	0.0282	0.0937	-1.48	-0.0152	0.8135	0.0135	0.4472

Figure Legends

Fig. 1. Sampling locations for *Entelurus aequoreus* in the northeastern Atlantic Ocean. Predominant ocean currents are shown in grey (after OSPAR, 2010). See Table 1 for sample information and abbreviations.

Fig. 2. Maximum likelihood networks for *Entelurus aequoreus* pooled over years for a) mtDNA *cytochrome b* sequences (900bp) and b) mtDNA control region sequences with microsatellite at position 287 removed (383bp). Each circle represents a haplotype and its size is proportional to its total frequency. Colors correspond to regions (see Table 1 and text for descriptions). Black crossbars represent a single nucleotide mutation and filled in circles represent reconstructed haplotypes not sampled.

Fig. 3. Bayesian Skyline Plot (BSP) showing changes in effective population size through time (thousand years before present, ka). Dashed line represents the median posterior estimate of the effective population size. The grey area delimited by continuous black lines shows the 95% highest posterior density limits.